

## 第2章 その他のデータ

基本健康診査

市民の健康状態

福生市健康関連事業の御案内

アンケート以外の健康に関する重要なデータをまとめました。




## 基本健康診査結果

福生市では、40歳以上の方を対象に基本健康診査を実施しています。ここでは、平成17年度に実施しました基本健康診査の結果を各検査項目ごとに「異常なし」「要経過観察」「要指導」「要医療」で表しています。


### 総合判定

| 年齢     | 性別 | 受診者    | 異常なし  |       | 要経過観察 |       | 要指導   |       | 要医療   |       |
|--------|----|--------|-------|-------|-------|-------|-------|-------|-------|-------|
|        |    |        | 人     | 率     | 人     | 率     | 人     | 率     | 人     | 率     |
| 75歳～   | 男  | 732    | 22    | 3.0%  | 195   | 26.6% | 54    | 7.4%  | 461   | 63.0% |
|        | 女  | 1,240  | 46    | 3.7%  | 285   | 23.0% | 101   | 8.1%  | 808   | 65.2% |
|        | 計  | 1,972  | 68    | 3.4%  | 480   | 24.3% | 155   | 7.9%  | 1269  | 64.4% |
| 70～75歳 | 男  | 621    | 24    | 3.9%  | 160   | 25.8% | 63    | 10.1% | 374   | 60.2% |
|        | 女  | 936    | 70    | 7.5%  | 236   | 25.2% | 101   | 10.8% | 529   | 56.5% |
|        | 計  | 1,557  | 94    | 6.1%  | 396   | 25.4% | 164   | 10.5% | 903   | 58.0% |
| 65～70歳 | 男  | 677    | 57    | 8.4%  | 155   | 22.9% | 82    | 12.1% | 383   | 56.6% |
|        | 女  | 940    | 86    | 9.2%  | 254   | 27.0% | 91    | 9.7%  | 509   | 54.1% |
|        | 計  | 1,617  | 143   | 8.8%  | 409   | 25.3% | 173   | 10.7% | 892   | 55.2% |
| 60～65歳 | 男  | 668    | 63    | 9.4%  | 157   | 23.5% | 80    | 12.0% | 368   | 55.1% |
|        | 女  | 999    | 123   | 12.3% | 246   | 24.6% | 135   | 13.5% | 495   | 49.6% |
|        | 計  | 1,667  | 186   | 11.1% | 403   | 24.2% | 215   | 12.9% | 863   | 51.8% |
| 55～60歳 | 男  | 395    | 42    | 10.6% | 99    | 25.1% | 46    | 11.6% | 208   | 52.7% |
|        | 女  | 841    | 123   | 14.6% | 240   | 28.5% | 103   | 12.3% | 375   | 44.6% |
|        | 計  | 1,236  | 165   | 13.3% | 339   | 27.4% | 149   | 12.1% | 583   | 47.2% |
| 50～55歳 | 男  | 280    | 34    | 12.1% | 71    | 25.4% | 39    | 13.9% | 136   | 48.6% |
|        | 女  | 720    | 143   | 19.9% | 245   | 34.0% | 76    | 10.6% | 256   | 35.5% |
|        | 計  | 1,000  | 177   | 17.7% | 316   | 31.6% | 115   | 11.5% | 392   | 39.2% |
| 45～50歳 | 男  | 255    | 39    | 15.3% | 78    | 30.6% | 39    | 15.3% | 99    | 38.8% |
|        | 女  | 546    | 155   | 28.4% | 167   | 30.6% | 60    | 11.0% | 164   | 30.0% |
|        | 計  | 801    | 194   | 24.2% | 245   | 30.6% | 99    | 12.4% | 263   | 32.8% |
| 40～45歳 | 男  | 230    | 41    | 17.8% | 78    | 33.9% | 26    | 11.3% | 85    | 37.0% |
|        | 女  | 563    | 170   | 30.2% | 196   | 34.8% | 60    | 10.7% | 137   | 24.3% |
|        | 計  | 793    | 211   | 26.6% | 274   | 34.6% | 86    | 10.8% | 222   | 28.0% |
| 不明     |    | 2      |       |       |       |       |       |       |       |       |
| 合計     | 男  | 3,858  | 322   | 8.4%  | 993   | 25.7% | 429   | 11.1% | 2,114 | 54.8% |
|        | 女  | 6,785  | 916   | 13.5% | 1,869 | 27.6% | 727   | 10.7% | 3,273 | 48.2% |
|        | 計  | 10,645 | 1,240 | 11.6% | 2,862 | 26.9% | 1,156 | 10.9% | 5,387 | 50.6% |




## 高血圧

| 年齢     | 性別 | 受診者    | 異常なし  |       | 要経過観察 |       | 要指導 |      | 要医療   |       |
|--------|----|--------|-------|-------|-------|-------|-----|------|-------|-------|
|        |    |        | 人     | 率     | 人     | 率     | 人   | 率    | 人     | 率     |
| 75歳～   | 男  | 732    | 406   | 55.5% | 74    | 10.1% | 5   | 0.7% | 247   | 33.7% |
|        | 女  | 1,240  | 606   | 48.9% | 133   | 10.7% | 18  | 1.4% | 483   | 39.0% |
|        | 計  | 1,972  | 1,012 | 51.3% | 207   | 10.5% | 23  | 1.2% | 730   | 37.0% |
| 70～75歳 | 男  | 621    | 302   | 48.6% | 84    | 13.5% | 13  | 2.1% | 222   | 35.8% |
|        | 女  | 936    | 508   | 54.3% | 110   | 11.8% | 20  | 2.1% | 298   | 31.8% |
|        | 計  | 1,557  | 810   | 52.0% | 194   | 12.5% | 33  | 2.1% | 520   | 33.4% |
| 65～70歳 | 男  | 677    | 389   | 57.4% | 81    | 12.0% | 12  | 1.8% | 195   | 28.8% |
|        | 女  | 940    | 557   | 59.3% | 104   | 11.1% | 20  | 2.1% | 259   | 27.5% |
|        | 計  | 1,617  | 946   | 58.5% | 185   | 11.4% | 32  | 2.0% | 454   | 28.1% |
| 60～65歳 | 男  | 668    | 381   | 57.0% | 72    | 10.8% | 29  | 4.3% | 186   | 27.9% |
|        | 女  | 999    | 658   | 65.9% | 92    | 9.2%  | 28  | 2.8% | 221   | 22.1% |
|        | 計  | 1,667  | 1,039 | 62.4% | 164   | 9.8%  | 57  | 3.4% | 407   | 24.4% |
| 55～60歳 | 男  | 395    | 233   | 59.0% | 49    | 12.4% | 9   | 2.3% | 104   | 26.3% |
|        | 女  | 841    | 617   | 73.4% | 71    | 8.4%  | 16  | 1.9% | 137   | 16.3% |
|        | 計  | 1,236  | 850   | 68.8% | 120   | 9.7%  | 25  | 2.0% | 241   | 19.5% |
| 50～55歳 | 男  | 280    | 188   | 67.2% | 32    | 11.4% | 2   | 0.7% | 58    | 20.7% |
|        | 女  | 720    | 582   | 80.8% | 41    | 5.7%  | 7   | 1.0% | 90    | 12.5% |
|        | 計  | 1,000  | 770   | 77.0% | 73    | 7.3%  | 9   | 0.9% | 148   | 14.8% |
| 45～50歳 | 男  | 255    | 194   | 76.1% | 27    | 10.6% | 3   | 1.2% | 31    | 12.1% |
|        | 女  | 546    | 470   | 86.1% | 34    | 6.2%  | 7   | 1.3% | 35    | 6.4%  |
|        | 計  | 801    | 664   | 82.9% | 61    | 7.6%  | 10  | 1.3% | 66    | 8.2%  |
| 40～45歳 | 男  | 230    | 184   | 80.0% | 20    | 8.7%  | 1   | 0.4% | 25    | 10.9% |
|        | 女  | 563    | 516   | 91.6% | 24    | 4.3%  | 2   | 0.4% | 21    | 3.7%  |
|        | 計  | 793    | 700   | 88.3% | 44    | 5.5%  | 3   | 0.4% | 46    | 5.8%  |
| 不明     |    | 2      |       |       |       |       |     |      |       |       |
| 合計     | 男  | 3,858  | 2,277 | 59.0% | 439   | 11.4% | 74  | 1.9% | 1,068 | 27.7% |
|        | 女  | 6,785  | 4,514 | 66.5% | 609   | 9.0%  | 118 | 1.7% | 1,544 | 22.8% |
|        | 計  | 10,645 | 6,793 | 63.8% | 1,048 | 9.9%  | 192 | 1.8% | 2,612 | 24.5% |




## 心疾患

| 年齢     | 性別 | 受診者    | 異常なし  |       | 要経過観察 |       | 要指導 |       | 要医療 |       |
|--------|----|--------|-------|-------|-------|-------|-----|-------|-----|-------|
|        |    |        | 人     | 率     | 人     | 率     | 人   | 率     | 人   | 率     |
| 75歳～   | 男  | 732    | 331   | 45.2% | 184   | 25.1% | 79  | 10.8% | 138 | 18.9% |
|        | 女  | 1,240  | 606   | 48.9% | 304   | 24.5% | 112 | 9.0%  | 218 | 17.6% |
|        | 計  | 1,972  | 937   | 47.5% | 488   | 24.7% | 191 | 9.7%  | 356 | 18.1% |
| 70～75歳 | 男  | 621    | 365   | 58.8% | 123   | 19.8% | 51  | 8.2%  | 82  | 13.2% |
|        | 女  | 936    | 561   | 59.9% | 198   | 21.2% | 77  | 8.2%  | 100 | 10.7% |
|        | 計  | 1,557  | 926   | 59.5% | 321   | 20.6% | 128 | 8.2%  | 182 | 11.7% |
| 65～70歳 | 男  | 677    | 468   | 69.1% | 104   | 15.4% | 41  | 6.0%  | 64  | 9.5%  |
|        | 女  | 940    | 632   | 67.2% | 178   | 19.0% | 66  | 7.0%  | 64  | 6.8%  |
|        | 計  | 1,617  | 1,100 | 68.0% | 282   | 17.5% | 107 | 6.6%  | 128 | 7.9%  |
| 60～65歳 | 男  | 668    | 479   | 71.7% | 103   | 15.4% | 36  | 5.4%  | 50  | 7.5%  |
|        | 女  | 999    | 760   | 76.1% | 139   | 13.9% | 57  | 5.7%  | 43  | 4.3%  |
|        | 計  | 1,667  | 1,239 | 74.3% | 242   | 14.5% | 93  | 5.6%  | 93  | 5.6%  |
| 55～60歳 | 男  | 395    | 277   | 70.1% | 74    | 18.7% | 20  | 5.1%  | 24  | 6.1%  |
|        | 女  | 841    | 684   | 81.2% | 105   | 12.5% | 25  | 3.0%  | 27  | 3.2%  |
|        | 計  | 1,236  | 961   | 77.8% | 179   | 14.5% | 45  | 3.6%  | 51  | 4.1%  |
| 50～55歳 | 男  | 280    | 208   | 74.3% | 48    | 17.1% | 14  | 5.0%  | 10  | 3.6%  |
|        | 女  | 720    | 585   | 81.2% | 99    | 13.8% | 20  | 2.8%  | 16  | 2.2%  |
|        | 計  | 1,000  | 793   | 79.3% | 147   | 14.7% | 34  | 3.4%  | 26  | 2.6%  |
| 45～50歳 | 男  | 255    | 211   | 82.8% | 28    | 11.0% | 8   | 3.1%  | 8   | 3.1%  |
|        | 女  | 546    | 458   | 83.9% | 64    | 11.7% | 15  | 2.7%  | 9   | 1.7%  |
|        | 計  | 801    | 669   | 83.5% | 92    | 11.5% | 23  | 2.9%  | 17  | 2.1%  |
| 40～45歳 | 男  | 230    | 185   | 80.4% | 35    | 15.2% | 5   | 2.2%  | 5   | 2.2%  |
|        | 女  | 563    | 467   | 83.0% | 80    | 14.2% | 8   | 1.4%  | 8   | 1.4%  |
|        | 計  | 793    | 652   | 82.2% | 115   | 14.6% | 13  | 1.6%  | 13  | 1.6%  |
| 不明     |    | 2      |       |       |       |       |     |       |     |       |
| 合計     | 男  | 3,858  | 2,524 | 65.4% | 699   | 18.1% | 254 | 6.6%  | 381 | 9.9%  |
|        | 女  | 6,785  | 4,753 | 70.1% | 1,167 | 17.2% | 380 | 5.6%  | 485 | 7.1%  |
|        | 計  | 10,645 | 7,279 | 68.4% | 1,866 | 17.5% | 634 | 6.0%  | 866 | 8.1%  |




## 動脈硬化

| 年齢     | 性別 | 受診者    | 異常なし  |       | 要経過観察 |       | 要指導 |      | 要医療 |      |
|--------|----|--------|-------|-------|-------|-------|-----|------|-----|------|
|        |    |        | 人     | 率     | 人     | 率     | 人   | 率    | 人   | 率    |
| 75歳～   | 男  | 732    | 617   | 84.3% | 83    | 11.3% | 11  | 1.5% | 21  | 2.9% |
|        | 女  | 1,240  | 1,017 | 82.0% | 160   | 12.9% | 6   | 0.5% | 57  | 4.6% |
|        | 計  | 1,972  | 1,634 | 82.9% | 243   | 12.3% | 17  | 0.9% | 78  | 3.9% |
| 70～75歳 | 男  | 621    | 551   | 88.7% | 52    | 8.4%  | 6   | 1.0% | 12  | 1.9% |
|        | 女  | 936    | 792   | 84.6% | 106   | 11.3% | 9   | 1.0% | 29  | 3.1% |
|        | 計  | 1,557  | 1,343 | 86.3% | 158   | 10.1% | 15  | 1.0% | 41  | 2.6% |
| 65～70歳 | 男  | 677    | 618   | 91.3% | 42    | 6.2%  | 2   | 0.3% | 15  | 2.2% |
|        | 女  | 940    | 847   | 90.1% | 69    | 7.4%  | 4   | 0.4% | 20  | 2.1% |
|        | 計  | 1,617  | 1,465 | 90.6% | 111   | 6.9%  | 6   | 0.4% | 35  | 2.1% |
| 60～65歳 | 男  | 668    | 624   | 93.4% | 30    | 4.5%  | 5   | 0.8% | 9   | 1.3% |
|        | 女  | 999    | 918   | 91.9% | 54    | 5.4%  | 8   | 0.8% | 19  | 1.9% |
|        | 計  | 1,667  | 1,542 | 92.5% | 84    | 5.0%  | 13  | 0.8% | 28  | 1.7% |
| 55～60歳 | 男  | 395    | 368   | 93.2% | 7     | 1.8%  | 4   | 1.0% | 16  | 4.0% |
|        | 女  | 841    | 792   | 94.2% | 36    | 4.3%  | 5   | 0.6% | 8   | 0.9% |
|        | 計  | 1,236  | 1,160 | 93.9% | 43    | 3.5%  | 9   | 0.7% | 24  | 1.9% |
| 50～55歳 | 男  | 280    | 270   | 96.4% | 3     | 1.1%  | 0   | 0.0% | 7   | 2.5% |
|        | 女  | 720    | 699   | 97.1% | 10    | 1.4%  | 2   | 0.3% | 9   | 1.2% |
|        | 計  | 1,000  | 969   | 96.9% | 13    | 1.3%  | 2   | 0.2% | 16  | 1.6% |
| 45～50歳 | 男  | 255    | 249   | 97.6% | 2     | 0.8%  | 3   | 1.2% | 1   | 0.4% |
|        | 女  | 546    | 537   | 98.3% | 7     | 1.3%  | 1   | 0.2% | 1   | 0.2% |
|        | 計  | 801    | 786   | 98.1% | 9     | 1.1%  | 4   | 0.5% | 2   | 0.3% |
| 40～45歳 | 男  | 230    | 225   | 97.8% | 0     | 0.0%  | 0   | 0.0% | 5   | 2.2% |
|        | 女  | 563    | 558   | 99.1% | 3     | 0.5%  | 0   | 0.0% | 2   | 0.4% |
|        | 計  | 793    | 783   | 98.7% | 3     | 0.4%  | 0   | 0.0% | 7   | 0.9% |
| 不明     |    | 2      |       |       |       |       |     |      |     |      |
| 合計     | 男  | 3,858  | 3,522 | 91.3% | 219   | 5.7%  | 31  | 0.8% | 86  | 2.2% |
|        | 女  | 6,785  | 6,160 | 90.8% | 445   | 6.6%  | 35  | 0.5% | 145 | 2.1% |
|        | 計  | 10,645 | 9,684 | 91.0% | 664   | 6.2%  | 66  | 0.6% | 231 | 2.2% |




## 高脂血症

| 年齢     | 性別 | 受診者    | 異常なし  |       | 要経過観察 |       | 要指導 |       | 要医療   |       |
|--------|----|--------|-------|-------|-------|-------|-----|-------|-------|-------|
|        |    |        | 人     | 率     | 人     | 率     | 人   | 率     | 人     | 率     |
| 75歳～   | 男  | 732    | 503   | 68.7% | 74    | 10.1% | 30  | 4.1%  | 125   | 17.1% |
|        | 女  | 1,240  | 654   | 52.7% | 180   | 14.5% | 80  | 6.5%  | 326   | 26.3% |
|        | 計  | 1,972  | 1,157 | 58.7% | 254   | 12.9% | 110 | 5.6%  | 451   | 22.8% |
| 70～75歳 | 男  | 621    | 363   | 58.5% | 100   | 16.1% | 38  | 6.1%  | 120   | 19.3% |
|        | 女  | 936    | 454   | 48.5% | 150   | 16.0% | 74  | 7.9%  | 258   | 27.6% |
|        | 計  | 1,557  | 817   | 52.5% | 250   | 16.0% | 112 | 7.2%  | 378   | 24.3% |
| 65～70歳 | 男  | 677    | 400   | 59.1% | 91    | 13.4% | 50  | 7.4%  | 136   | 20.1% |
|        | 女  | 940    | 415   | 44.1% | 182   | 19.4% | 73  | 7.8%  | 270   | 28.7% |
|        | 計  | 1,617  | 815   | 50.4% | 273   | 16.9% | 123 | 7.6%  | 406   | 25.1% |
| 60～65歳 | 男  | 668    | 375   | 56.1% | 109   | 16.3% | 56  | 8.4%  | 128   | 19.2% |
|        | 女  | 999    | 433   | 43.4% | 191   | 19.1% | 106 | 10.6% | 269   | 26.9% |
|        | 計  | 1,667  | 808   | 48.5% | 300   | 18.0% | 162 | 9.7%  | 397   | 23.8% |
| 55～60歳 | 男  | 395    | 196   | 49.6% | 69    | 17.5% | 37  | 9.4%  | 93    | 23.5% |
|        | 女  | 841    | 389   | 46.2% | 151   | 18.0% | 76  | 9.0%  | 225   | 26.8% |
|        | 計  | 1,236  | 585   | 47.3% | 220   | 17.8% | 113 | 9.2%  | 318   | 25.7% |
| 50～55歳 | 男  | 280    | 132   | 47.2% | 44    | 15.7% | 25  | 8.9%  | 79    | 28.2% |
|        | 女  | 720    | 411   | 57.1% | 129   | 17.9% | 56  | 7.8%  | 124   | 17.2% |
|        | 計  | 1,000  | 543   | 54.3% | 173   | 17.3% | 81  | 8.1%  | 203   | 20.3% |
| 45～50歳 | 男  | 255    | 135   | 52.9% | 43    | 16.9% | 23  | 9.0%  | 54    | 21.2% |
|        | 女  | 546    | 390   | 71.4% | 79    | 14.5% | 31  | 5.7%  | 46    | 8.4%  |
|        | 計  | 801    | 525   | 65.6% | 122   | 15.2% | 54  | 6.7%  | 100   | 12.5% |
| 40～45歳 | 男  | 230    | 118   | 51.3% | 39    | 17.0% | 18  | 7.8%  | 55    | 23.9% |
|        | 女  | 563    | 439   | 78.0% | 74    | 13.1% | 27  | 4.8%  | 23    | 4.1%  |
|        | 計  | 793    | 557   | 70.2% | 113   | 14.3% | 45  | 5.7%  | 78    | 9.8%  |
| 不明     |    | 2      |       |       |       |       |     |       |       |       |
| 合計     | 男  | 3,858  | 2,222 | 57.6% | 569   | 14.7% | 277 | 7.2%  | 790   | 20.5% |
|        | 女  | 6,785  | 3,585 | 52.8% | 1,136 | 16.8% | 523 | 7.7%  | 1,541 | 22.7% |
|        | 計  | 10,645 | 5,809 | 54.6% | 1,705 | 16.0% | 800 | 7.5%  | 2,331 | 21.9% |



## 呼吸器疾患


| 年齢     | 性別 | 受診者    | 異常なし  |       | 要経過観察 |       | 要指導 |      | 要医療 |      |
|--------|----|--------|-------|-------|-------|-------|-----|------|-----|------|
|        |    |        | 人     | 率     | 人     | 率     | 人   | 率    | 人   | 率    |
| 75歳～   | 男  | 732    | 479   | 65.4% | 189   | 25.8% | 7   | 1.0% | 57  | 7.8% |
|        | 女  | 1,240  | 943   | 76.1% | 241   | 19.4% | 6   | 0.5% | 50  | 4.0% |
|        | 計  | 1,972  | 1,422 | 72.1% | 430   | 21.8% | 13  | 0.7% | 107 | 5.4% |
| 70～75歳 | 男  | 621    | 460   | 74.1% | 124   | 19.9% | 8   | 1.3% | 29  | 4.7% |
|        | 女  | 936    | 779   | 83.2% | 119   | 12.7% | 9   | 1.0% | 29  | 3.1% |
|        | 計  | 1,557  | 1,239 | 79.6% | 243   | 15.6% | 17  | 1.1% | 58  | 3.7% |
| 65～70歳 | 男  | 677    | 549   | 81.1% | 109   | 16.1% | 5   | 0.7% | 14  | 2.1% |
|        | 女  | 940    | 809   | 86.1% | 110   | 11.7% | 3   | 0.3% | 18  | 1.9% |
|        | 計  | 1,617  | 1,358 | 84.0% | 219   | 13.5% | 8   | 0.5% | 32  | 2.0% |
| 60～65歳 | 男  | 668    | 558   | 83.5% | 82    | 12.3% | 6   | 0.9% | 22  | 3.3% |
|        | 女  | 999    | 893   | 89.4% | 78    | 7.8%  | 7   | 0.7% | 21  | 2.1% |
|        | 計  | 1,667  | 1,451 | 87.0% | 160   | 9.6%  | 13  | 0.8% | 43  | 2.6% |
| 55～60歳 | 男  | 395    | 338   | 85.6% | 48    | 12.1% | 4   | 1.0% | 5   | 1.3% |
|        | 女  | 841    | 775   | 92.2% | 55    | 6.5%  | 4   | 0.5% | 7   | 0.8% |
|        | 計  | 1,236  | 1,113 | 90.0% | 103   | 8.3%  | 8   | 0.7% | 12  | 1.0% |
| 50～55歳 | 男  | 280    | 245   | 87.5% | 25    | 8.9%  | 3   | 1.1% | 7   | 2.5% |
|        | 女  | 720    | 662   | 91.9% | 43    | 6.0%  | 5   | 0.7% | 10  | 1.4% |
|        | 計  | 1,000  | 907   | 90.7% | 68    | 6.8%  | 8   | 0.8% | 17  | 1.7% |
| 45～50歳 | 男  | 255    | 229   | 89.8% | 22    | 8.6%  | 0   | 0.0% | 4   | 1.6% |
|        | 女  | 546    | 512   | 93.8% | 23    | 4.2%  | 2   | 0.4% | 9   | 1.6% |
|        | 計  | 801    | 741   | 92.5% | 45    | 5.6%  | 2   | 0.3% | 13  | 1.6% |
| 40～45歳 | 男  | 230    | 213   | 92.6% | 15    | 6.5%  | 0   | 0.0% | 2   | 0.9% |
|        | 女  | 563    | 523   | 92.9% | 32    | 5.7%  | 3   | 0.5% | 5   | 0.9% |
|        | 計  | 793    | 736   | 92.8% | 47    | 5.9%  | 3   | 0.4% | 7   | 0.9% |
| 不明     |    | 2      |       |       |       |       |     |      |     |      |
| 合計     | 男  | 3,858  | 3,071 | 79.6% | 614   | 15.9% | 33  | 0.9% | 140 | 3.6% |
|        | 女  | 6,785  | 5,896 | 86.9% | 701   | 10.3% | 39  | 0.6% | 149 | 2.2% |
|        | 計  | 10,645 | 8,969 | 84.3% | 1,315 | 12.3% | 72  | 0.7% | 289 | 2.7% |






## 糖尿病

| 年齢     | 性別 | 受診者    | 異常なし  |       | 要経過観察 |      | 要指導 |       | 要医療   |       |
|--------|----|--------|-------|-------|-------|------|-----|-------|-------|-------|
|        |    |        | 人     | 率     | 人     | 率    | 人   | 率     | 人     | 率     |
| 75歳～   | 男  | 732    | 536   | 73.2% | 71    | 9.7% | 13  | 1.8%  | 112   | 15.3% |
|        | 女  | 1,240  | 972   | 78.4% | 95    | 7.7% | 16  | 1.3%  | 157   | 12.6% |
|        | 計  | 1,972  | 1,508 | 76.5% | 166   | 8.4% | 29  | 1.5%  | 269   | 13.6% |
| 70～75歳 | 男  | 621    | 459   | 73.9% | 48    | 7.7% | 14  | 2.3%  | 100   | 16.1% |
|        | 女  | 936    | 759   | 81.1% | 58    | 6.2% | 9   | 1.0%  | 110   | 11.7% |
|        | 計  | 1,557  | 1,218 | 78.2% | 106   | 6.8% | 23  | 1.5%  | 210   | 13.5% |
| 65～70歳 | 男  | 677    | 511   | 75.5% | 46    | 6.8% | 8   | 1.2%  | 112   | 16.5% |
|        | 女  | 940    | 774   | 82.4% | 53    | 5.6% | 17  | 1.8%  | 96    | 10.2% |
|        | 計  | 1,617  | 1,285 | 79.5% | 99    | 6.1% | 25  | 1.5%  | 208   | 12.9% |
| 60～65歳 | 男  | 668    | 523   | 78.3% | 42    | 6.3% | 10  | 1.5%  | 93    | 13.9% |
|        | 女  | 999    | 854   | 85.5% | 49    | 4.9% | 9   | 0.9%  | 87    | 8.7%  |
|        | 計  | 1,667  | 1,377 | 82.6% | 91    | 5.5% | 19  | 1.1%  | 180   | 10.8% |
| 55～60歳 | 男  | 395    | 309   | 78.2% | 29    | 7.4% | 6   | 1.5%  | 51    | 12.9% |
|        | 女  | 841    | 745   | 88.6% | 29    | 3.5% | 7   | 0.8%  | 60    | 7.1%  |
|        | 計  | 1,236  | 1,054 | 85.3% | 58    | 4.7% | 13  | 1.0%  | 111   | 9.0%  |
| 50～55歳 | 男  | 280    | 225   | 80.4% | 18    | 6.4% | 2   | 0.7%  | 35    | 12.5% |
|        | 女  | 720    | 676   | 93.9% | 11    | 1.5% | 5   | 0.7%  | 28    | 3.9%  |
|        | 計  | 1,000  | 901   | 90.1% | 29    | 2.9% | 7   | 0.7%  | 63    | 6.3%  |
| 45～50歳 | 男  | 255    | 230   | 90.2% | 12    | 4.7% | 1   | 0.4%  | 12    | 4.7%  |
|        | 女  | 546    | 521   | 95.4% | 11    | 2.0% | 1   | 0.2%  | 13    | 2.4%  |
|        | 計  | 801    | 751   | 93.8% | 23    | 2.9% | 2   | 0.2%  | 25    | 3.1%  |
| 40～45歳 | 男  | 230    | 217   | 94.4% | 4     | 1.7% | 3   | 1.3%  | 6     | 2.6%  |
|        | 女  | 563    | 543   | 96.5% | 7     | 1.2% | 0   | 0.0%  | 13    | 2.3%  |
|        | 計  | 793    | 760   | 95.8% | 11    | 1.4% | 3   | 0.4%  | 19    | 2.4%  |
| 不明     |    | 2      |       |       |       |      |     |       |       |       |
| 合計     | 男  | 3,858  | 3,010 | 78.0% | 270   | 7.0% | 57  | 1.5%  | 521   | 13.5% |
|        | 女  | 6,785  | 5,844 | 86.1% | 313   | 4.6% | 64  | 10.0% | 564   | 8.3%  |
|        | 計  | 10,645 | 8,856 | 83.2% | 583   | 5.5% | 121 | 1.1%  | 1,085 | 10.2% |




# 胃腸病

| 年齢     | 性別 | 受診者    | 異常なし   |       | 要経過観察 |      | 要指導 |      | 要医療 |      |
|--------|----|--------|--------|-------|-------|------|-----|------|-----|------|
|        |    |        | 人      | 率     | 人     | 率    | 人   | 率    | 人   | 率    |
| 75歳～   | 男  | 732    | 659    | 90.0% | 30    | 4.1% | 9   | 1.2% | 34  | 4.7% |
|        | 女  | 1,240  | 1,138  | 91.8% | 36    | 2.9% | 14  | 1.1% | 52  | 4.2% |
|        | 計  | 1,972  | 1,797  | 91.1% | 66    | 3.3% | 23  | 1.2% | 86  | 4.4% |
| 70～75歳 | 男  | 621    | 569    | 91.6% | 19    | 3.1% | 8   | 1.3% | 25  | 4.0% |
|        | 女  | 936    | 876    | 93.6% | 14    | 1.5% | 12  | 1.3% | 34  | 3.6% |
|        | 計  | 1,557  | 1,445  | 92.8% | 33    | 2.1% | 20  | 1.3% | 59  | 3.8% |
| 65～70歳 | 男  | 677    | 642    | 94.8% | 7     | 1.0% | 4   | 0.6% | 24  | 3.6% |
|        | 女  | 940    | 896    | 95.3% | 14    | 1.5% | 5   | 0.5% | 25  | 2.7% |
|        | 計  | 1,617  | 1,538  | 95.1% | 21    | 1.3% | 9   | 0.6% | 49  | 3.0% |
| 60～65歳 | 男  | 668    | 623    | 93.3% | 12    | 1.8% | 5   | 0.7% | 28  | 4.2% |
|        | 女  | 999    | 955    | 95.6% | 14    | 1.4% | 6   | 0.6% | 24  | 2.4% |
|        | 計  | 1,667  | 1,578  | 94.7% | 26    | 1.5% | 11  | 0.7% | 52  | 3.1% |
| 55～60歳 | 男  | 395    | 376    | 95.2% | 4     | 1.0% | 0   | 0.0% | 15  | 3.8% |
|        | 女  | 841    | 819    | 97.4% | 6     | 0.7% | 4   | 0.5% | 12  | 1.4% |
|        | 計  | 1,236  | 1,195  | 96.7% | 10    | 0.8% | 4   | 0.3% | 27  | 2.2% |
| 50～55歳 | 男  | 280    | 273    | 97.5% | 1     | 0.3% | 3   | 1.1% | 3   | 1.1% |
|        | 女  | 720    | 696    | 96.7% | 4     | 0.5% | 3   | 0.4% | 17  | 2.4% |
|        | 計  | 1,000  | 969    | 96.9% | 5     | 0.5% | 6   | 0.6% | 20  | 2.0% |
| 45～50歳 | 男  | 255    | 243    | 95.3% | 4     | 1.6% | 1   | 0.4% | 7   | 2.7% |
|        | 女  | 546    | 527    | 96.5% | 7     | 1.3% | 2   | 0.4% | 10  | 1.8% |
|        | 計  | 801    | 770    | 96.1% | 11    | 1.4% | 3   | 0.4% | 17  | 2.1% |
| 40～45歳 | 男  | 230    | 220    | 95.7% | 6     | 2.6% | 1   | 0.4% | 3   | 1.3% |
|        | 女  | 563    | 540    | 95.9% | 6     | 1.1% | 5   | 0.9% | 12  | 2.1% |
|        | 計  | 793    | 760    | 95.8% | 12    | 1.5% | 6   | 0.8% | 15  | 1.9% |
| 不明     |    | 2      |        |       |       |      |     |      |     |      |
| 合計     | 男  | 3,858  | 3,605  | 93.4% | 83    | 2.2% | 31  | 0.8% | 139 | 3.6% |
|        | 女  | 6,785  | 6,447  | 95.0% | 101   | 1.5% | 51  | 0.8% | 186 | 2.7% |
|        | 計  | 10,645 | 10,054 | 94.4% | 184   | 1.7% | 82  | 0.8% | 325 | 3.1% |




## 肝疾患

| 年齢     | 性別 | 受診者    | 異常なし  |       | 要経過観察 |       | 要指導 |      | 要医療 |       |
|--------|----|--------|-------|-------|-------|-------|-----|------|-----|-------|
|        |    |        | 人     | 率     | 人     | 率     | 人   | 率    | 人   | 率     |
| 75歳～   | 男  | 732    | 593   | 81.0% | 76    | 10.4% | 21  | 2.9% | 42  | 5.7%  |
|        | 女  | 1,240  | 1,053 | 84.9% | 120   | 9.7%  | 25  | 2.0% | 42  | 3.4%  |
|        | 計  | 1,972  | 1,646 | 83.5% | 196   | 9.9%  | 46  | 2.3% | 84  | 4.3%  |
| 70～75歳 | 男  | 621    | 497   | 80.0% | 67    | 10.8% | 17  | 2.7% | 40  | 6.5%  |
|        | 女  | 936    | 792   | 84.6% | 88    | 9.4%  | 20  | 2.1% | 36  | 3.9%  |
|        | 計  | 1,557  | 1,289 | 82.8% | 155   | 9.9%  | 37  | 2.4% | 76  | 4.9%  |
| 65～70歳 | 男  | 677    | 500   | 73.9% | 82    | 12.1% | 40  | 5.9% | 55  | 8.1%  |
|        | 女  | 940    | 794   | 84.5% | 90    | 9.6%  | 23  | 2.4% | 33  | 3.5%  |
|        | 計  | 1,617  | 1,294 | 80.0% | 172   | 10.6% | 63  | 3.9% | 88  | 5.5%  |
| 60～65歳 | 男  | 668    | 517   | 77.4% | 79    | 11.8% | 28  | 4.2% | 44  | 6.6%  |
|        | 女  | 999    | 879   | 88.0% | 73    | 7.3%  | 9   | 0.9% | 38  | 3.8%  |
|        | 計  | 1,667  | 1,396 | 83.8% | 152   | 9.1%  | 37  | 2.2% | 82  | 4.9%  |
| 55～60歳 | 男  | 395    | 270   | 68.4% | 55    | 13.9% | 21  | 5.3% | 49  | 12.4% |
|        | 女  | 841    | 739   | 87.9% | 56    | 6.6%  | 21  | 2.5% | 25  | 3.0%  |
|        | 計  | 1,236  | 1,009 | 81.6% | 111   | 9.0%  | 42  | 3.4% | 74  | 6.0%  |
| 50～55歳 | 男  | 280    | 197   | 70.4% | 44    | 15.7% | 16  | 5.7% | 23  | 8.2%  |
|        | 女  | 720    | 630   | 87.5% | 64    | 8.9%  | 14  | 1.9% | 12  | 1.7%  |
|        | 計  | 1,000  | 827   | 82.7% | 108   | 10.8% | 30  | 3.0% | 35  | 3.5%  |
| 45～50歳 | 男  | 255    | 181   | 71.0% | 40    | 15.7% | 14  | 5.5% | 20  | 7.8%  |
|        | 女  | 546    | 497   | 91.0% | 32    | 5.9%  | 6   | 1.1% | 11  | 2.0%  |
|        | 計  | 801    | 678   | 84.6% | 72    | 9.0%  | 20  | 2.5% | 31  | 3.9%  |
| 40～45歳 | 男  | 230    | 162   | 70.4% | 39    | 17.0% | 10  | 4.3% | 19  | 8.3%  |
|        | 女  | 563    | 520   | 92.3% | 27    | 4.8%  | 6   | 1.1% | 10  | 1.8%  |
|        | 計  | 793    | 682   | 86.0% | 66    | 8.3%  | 16  | 2.0% | 29  | 3.7%  |
| 不明     |    | 2      |       |       |       |       |     |      |     |       |
| 合計     | 男  | 3,858  | 2,917 | 75.6% | 482   | 12.5% | 167 | 4.3% | 292 | 7.6%  |
|        | 女  | 6,785  | 5,904 | 87.0% | 550   | 8.1%  | 124 | 1.8% | 207 | 3.1%  |
|        | 計  | 10,645 | 8,823 | 82.9% | 1,032 | 9.7%  | 291 | 2.7% | 499 | 4.7%  |




# 腎疾患

| 年齢     | 性別 | 受診者    | 異常なし  |       | 要経過観察 |       | 要指導 |      | 要医療 |      |
|--------|----|--------|-------|-------|-------|-------|-----|------|-----|------|
|        |    |        | 人     | 率     | 人     | 率     | 人   | 率    | 人   | 率    |
| 75歳～   | 男  | 732    | 522   | 71.3% | 136   | 18.6% | 15  | 2.0% | 59  | 8.1% |
|        | 女  | 1,240  | 838   | 67.6% | 288   | 23.2% | 19  | 1.5% | 95  | 7.7% |
|        | 計  | 1,972  | 1,360 | 69.0% | 424   | 21.5% | 34  | 1.7% | 154 | 7.8% |
| 70～75歳 | 男  | 621    | 478   | 77.0% | 112   | 18.0% | 7   | 1.1% | 24  | 3.9% |
|        | 女  | 936    | 689   | 73.6% | 161   | 17.2% | 14  | 1.5% | 72  | 7.7% |
|        | 計  | 1,557  | 1,167 | 75.0% | 273   | 17.5% | 21  | 1.3% | 96  | 6.2% |
| 65～70歳 | 男  | 677    | 547   | 80.8% | 107   | 15.8% | 6   | 0.9% | 17  | 2.5% |
|        | 女  | 940    | 715   | 76.1% | 171   | 18.2% | 9   | 0.9% | 45  | 4.8% |
|        | 計  | 1,617  | 1,262 | 78.1% | 278   | 17.2% | 15  | 0.9% | 62  | 3.8% |
| 60～65歳 | 男  | 668    | 533   | 79.8% | 100   | 15.0% | 12  | 1.8% | 23  | 3.4% |
|        | 女  | 999    | 749   | 75.0% | 186   | 18.6% | 13  | 1.3% | 51  | 5.1% |
|        | 計  | 1,667  | 1,282 | 76.9% | 286   | 17.2% | 25  | 1.5% | 74  | 4.4% |
| 55～60歳 | 男  | 395    | 326   | 82.6% | 53    | 13.4% | 6   | 1.5% | 10  | 2.5% |
|        | 女  | 841    | 648   | 77.0% | 147   | 17.5% | 8   | 1.0% | 38  | 4.5% |
|        | 計  | 1,236  | 974   | 78.8% | 200   | 16.2% | 14  | 1.1% | 48  | 3.9% |
| 50～55歳 | 男  | 280    | 237   | 84.7% | 32    | 11.4% | 5   | 1.8% | 6   | 2.1% |
|        | 女  | 720    | 545   | 75.7% | 132   | 18.3% | 7   | 1.0% | 36  | 5.0% |
|        | 計  | 1,000  | 782   | 78.2% | 164   | 16.4% | 12  | 1.2% | 42  | 4.2% |
| 45～50歳 | 男  | 255    | 209   | 81.9% | 38    | 14.9% | 3   | 1.2% | 5   | 2.0% |
|        | 女  | 546    | 449   | 82.2% | 71    | 13.0% | 3   | 0.6% | 23  | 4.2% |
|        | 計  | 801    | 658   | 82.1% | 109   | 13.6% | 6   | 0.8% | 28  | 3.5% |
| 40～45歳 | 男  | 230    | 202   | 87.8% | 18    | 7.8%  | 1   | 0.5% | 9   | 3.9% |
|        | 女  | 563    | 438   | 77.8% | 86    | 15.3% | 4   | 0.7% | 35  | 6.2% |
|        | 計  | 793    | 640   | 80.7% | 104   | 13.1% | 5   | 0.6% | 44  | 5.6% |
| 不明     |    | 2      |       |       |       |       |     |      |     |      |
| 合計     | 男  | 3,858  | 3,054 | 79.2% | 596   | 15.4% | 55  | 1.4% | 153 | 4.0% |
|        | 女  | 6,785  | 5,071 | 74.8% | 1,242 | 18.3% | 77  | 1.1% | 395 | 5.8% |
|        | 計  | 10,645 | 8,127 | 76.3% | 1,838 | 17.3% | 132 | 1.2% | 548 | 5.2% |



## 血液疾患

| 年齢     | 性別 | 受診者    | 異常なし  |       | 要経過観察 |       | 要指導 |      | 要医療 |      |
|--------|----|--------|-------|-------|-------|-------|-----|------|-----|------|
|        |    |        | 人     | 率     | 人     | 率     | 人   | 率    | 人   | 率    |
| 75歳～   | 男  | 732    | 462   | 63.1% | 165   | 22.5% | 37  | 5.1% | 68  | 9.3% |
|        | 女  | 1,240  | 971   | 78.3% | 149   | 12.0% | 37  | 3.0% | 83  | 6.7% |
|        | 計  | 1,972  | 1,433 | 72.7% | 314   | 15.9% | 74  | 3.7% | 151 | 7.7% |
| 70～75歳 | 男  | 621    | 492   | 79.2% | 89    | 14.3% | 22  | 3.6% | 18  | 2.9% |
|        | 女  | 936    | 820   | 87.6% | 86    | 9.2%  | 15  | 1.6% | 15  | 1.6% |
|        | 計  | 1,557  | 1,312 | 84.3% | 175   | 11.2% | 37  | 2.4% | 33  | 2.1% |
| 65～70歳 | 男  | 677    | 577   | 85.2% | 59    | 8.7%  | 22  | 3.3% | 19  | 2.8% |
|        | 女  | 940    | 868   | 92.3% | 49    | 5.2%  | 7   | 0.8% | 16  | 1.7% |
|        | 計  | 1,617  | 1,445 | 89.3% | 108   | 6.7%  | 29  | 1.8% | 35  | 2.2% |
| 60～65歳 | 男  | 668    | 575   | 86.1% | 46    | 6.9%  | 25  | 3.7% | 22  | 3.3% |
|        | 女  | 999    | 940   | 94.1% | 36    | 3.6%  | 17  | 1.7% | 6   | 0.6% |
|        | 計  | 1,667  | 1,515 | 90.9% | 82    | 4.9%  | 42  | 2.5% | 28  | 1.7% |
| 55～60歳 | 男  | 395    | 345   | 87.4% | 27    | 6.8%  | 11  | 2.8% | 12  | 3.0% |
|        | 女  | 841    | 781   | 92.9% | 39    | 4.6%  | 14  | 1.7% | 7   | 0.8% |
|        | 計  | 1,236  | 1,126 | 91.1% | 66    | 5.3%  | 25  | 2.0% | 19  | 1.6% |
| 50～55歳 | 男  | 280    | 250   | 89.3% | 14    | 5.0%  | 6   | 2.1% | 10  | 3.6% |
|        | 女  | 720    | 640   | 88.9% | 50    | 6.9%  | 15  | 2.1% | 15  | 2.1% |
|        | 計  | 1,000  | 890   | 89.0% | 64    | 6.4%  | 21  | 2.1% | 25  | 2.5% |
| 45～50歳 | 男  | 255    | 237   | 92.9% | 11    | 4.3%  | 4   | 1.6% | 3   | 1.2% |
|        | 女  | 546    | 450   | 82.4% | 42    | 7.7%  | 16  | 2.9% | 38  | 7.0% |
|        | 計  | 801    | 687   | 85.8% | 53    | 6.6%  | 20  | 2.5% | 41  | 5.1% |
| 40～45歳 | 男  | 230    | 213   | 92.6% | 12    | 5.2%  | 1   | 0.4% | 4   | 1.8% |
|        | 女  | 563    | 464   | 82.4% | 38    | 6.7%  | 19  | 3.4% | 42  | 7.5% |
|        | 計  | 793    | 677   | 85.4% | 50    | 6.3%  | 20  | 2.5% | 46  | 5.8% |
| 不明     |    | 2      |       |       |       |       |     |      |     |      |
| 合計     | 男  | 3,858  | 3,151 | 81.7% | 423   | 11.0% | 128 | 3.3% | 156 | 4.0% |
|        | 女  | 6,785  | 5,934 | 87.4% | 489   | 7.2%  | 140 | 2.1% | 222 | 3.3% |
|        | 計  | 10,645 | 9,087 | 85.4% | 912   | 8.6%  | 268 | 2.5% | 378 | 3.5% |



## その他

| 年齢     | 性別 | 受診者    | 異常なし  |       | 要経過観察 |       | 要指導 |      | 要医療 |      |
|--------|----|--------|-------|-------|-------|-------|-----|------|-----|------|
|        |    |        | 人     | 率     | 人     | 率     | 人   | 率    | 人   | 率    |
| 75歳～   | 男  | 732    | 549   | 75.0% | 95    | 13.0% | 18  | 2.4% | 70  | 9.6% |
|        | 女  | 1,240  | 995   | 80.2% | 106   | 8.6%  | 42  | 3.4% | 97  | 7.8% |
|        | 計  | 1,972  | 1,544 | 78.3% | 201   | 10.2% | 60  | 3.0% | 167 | 8.5% |
| 70～75歳 | 男  | 621    | 501   | 80.7% | 54    | 8.7%  | 28  | 4.5% | 38  | 6.1% |
|        | 女  | 936    | 788   | 84.2% | 78    | 8.3%  | 23  | 2.5% | 47  | 5.0% |
|        | 計  | 1,557  | 1,289 | 82.8% | 132   | 8.5%  | 51  | 3.3% | 85  | 5.4% |
| 65～70歳 | 男  | 677    | 549   | 81.1% | 61    | 9.0%  | 26  | 3.8% | 41  | 6.1% |
|        | 女  | 940    | 819   | 87.1% | 67    | 7.1%  | 22  | 2.4% | 32  | 3.4% |
|        | 計  | 1,617  | 1,368 | 84.6% | 128   | 7.9%  | 48  | 3.0% | 73  | 4.5% |
| 60～65歳 | 男  | 668    | 566   | 84.7% | 44    | 6.6%  | 16  | 2.4% | 42  | 6.3% |
|        | 女  | 999    | 873   | 87.4% | 67    | 6.7%  | 27  | 2.7% | 32  | 3.2% |
|        | 計  | 1,667  | 1,439 | 86.3% | 111   | 6.7%  | 43  | 2.6% | 74  | 4.4% |
| 55～60歳 | 男  | 395    | 327   | 82.8% | 30    | 7.6%  | 12  | 3.0% | 26  | 6.6% |
|        | 女  | 841    | 748   | 89.0% | 55    | 6.5%  | 21  | 2.5% | 17  | 2.0% |
|        | 計  | 1,236  | 1,075 | 87.0% | 85    | 6.9%  | 33  | 2.6% | 43  | 3.5% |
| 50～55歳 | 男  | 280    | 229   | 81.8% | 16    | 5.7%  | 12  | 4.3% | 23  | 8.2% |
|        | 女  | 720    | 646   | 89.7% | 46    | 6.4%  | 19  | 2.6% | 9   | 1.3% |
|        | 計  | 1,000  | 875   | 87.5% | 62    | 6.2%  | 31  | 3.1% | 32  | 3.2% |
| 45～50歳 | 男  | 255    | 205   | 80.4% | 24    | 9.4%  | 9   | 3.5% | 17  | 6.7% |
|        | 女  | 546    | 491   | 89.9% | 27    | 5.0%  | 16  | 2.9% | 12  | 2.2% |
|        | 計  | 801    | 696   | 86.9% | 51    | 6.4%  | 25  | 3.1% | 29  | 3.6% |
| 40～45歳 | 男  | 230    | 178   | 77.4% | 20    | 8.7%  | 12  | 5.2% | 20  | 8.7% |
|        | 女  | 563    | 511   | 90.7% | 30    | 5.3%  | 11  | 2.0% | 11  | 2.0% |
|        | 計  | 793    | 689   | 86.9% | 50    | 6.3%  | 23  | 2.9% | 31  | 3.9% |
| 不明     |    | 2      |       |       |       |       |     |      |     |      |
| 年齢     | 性別 | 受診者    | 異常なし  |       | 要経過観察 |       | 要指導 |      | 要医療 |      |
|        |    |        | 人     | 率     | 人     | 率     | 人   | 率    | 人   | 率    |
| 合計     | 男  | 3,858  | 3,104 | 80.5% | 344   | 8.9%  | 133 | 3.4% | 277 | 7.2% |
|        | 女  | 6,785  | 5,871 | 86.5% | 476   | 7.0%  | 181 | 2.7% | 257 | 3.8% |
|        | 計  | 10,645 | 8,977 | 84.3% | 820   | 7.7%  | 314 | 3.0% | 534 | 5.0% |

